

# CH 221 Chapter Five Concept Guide

## 1. Specific Heat

### Question

How much heat is needed to raise the temperature of 28 g of aluminum from 27 °C to 163 °C? No phase changes occur. The specific heat of aluminum is 0.902 J/K\*g.

### Approach

The amount of heat needed is found from the known mass of Al, the change in temperature, and the specific heat of Al. The following equation relates heat to these values:  $q = (\text{mass})(\text{specific heat})(\text{temperature change})$ .

### Solution

$$\Delta T = 163\text{ }^{\circ}\text{C} - 27\text{ }^{\circ}\text{C} = 136\text{ }^{\circ}\text{C}, \text{ or } 136\text{ K.}$$

$$q = (\text{mass})(\text{specific heat})(\text{temperature change})$$

$$q = (28\text{ g})(0.902\text{ J/K}\cdot\text{g})(136\text{ K})$$

$$q = 3400\text{ J}$$

Note that the temperature increased and  $q$  is positive. Heat was added to the aluminum, so the direction of heat flow was into the system.

## 2. Specific Heat

### Question

What quantity of heat is needed to raise the temperature of 130 g of water by 5.0 °C? The specific heat of water is 4.184 J/K\*g.

### Approach

The following equation relates heat to mass and temperature change:

$$q = (\text{mass})(\text{specific heat})(\text{temperature change}).$$

### Solution

$$\Delta T = 5.0\text{ }^{\circ}\text{C}, \text{ or } 5.0\text{ K}$$

$$q = (\text{mass})(\text{specific heat})(\text{temperature change})$$

$$q = (130\text{ g})(4.184\text{ J/K}\cdot\text{g})(5.0\text{ K})$$

$$q = 2700\text{ J}$$

Note that the temperature increased and  $q$  is positive.

## 3. Molar Heat Capacity

### Problem

125 J of heat was absorbed by 1.00 mol of iron, resulting in a temperature change of 4.02 °C. Calculate the molar heat capacity of iron.

## Approach

Molar heat capacity can be calculated using the following equation:

$$q = (\text{number of moles})(\text{molar heat capacity})(\text{temperature change}).$$

## Solution

$$q = (\text{number of moles})(\text{molar heat capacity})(\text{temperature change}).$$

$$125 \text{ J} = (1.00 \text{ mol iron})(\text{molar heat capacity of iron})(4.02 \text{ }^\circ\text{C})$$

$$\text{molar heat capacity of iron} = 125 \text{ J} / (1.00 \text{ mol iron})(4.02 \text{ }^\circ\text{C})$$

$$\text{molar heat capacity of iron} = 31.1 \text{ J/mol}\cdot\text{K}$$

## 4. Molar Heat Capacity

Heat capacity values are tabulated in two ways: as molar heat capacity, and as specific heat capacity, which has already been discussed in this lesson. Molar heat capacity differs from specific heat capacity in that the former is the amount of heat required to raise the temperature of one mole of a substance by one Kelvin ( $\text{J/K}\cdot\text{g}$ ). You may recall that the specific heat is the amount of heat required to raise the temperature of one gram of a substance by one Kelvin ( $\text{J/K}\cdot\text{g}$ ).

The molar heat capacity may be calculated using the following equation:


$$q = (\text{number of moles})(\text{molar heat capacity})(\text{temperature change})$$

where  $q$  is heat in joules. The molar heat capacity has units of  $\text{J/K}\cdot\text{g}$ . For example, the specific heat of Cu is  $0.385 \text{ J/K}\cdot\text{g}$ . Its molar heat capacity is, therefore,  $24.5 \text{ J/K}\cdot\text{mol}$ .

## 5. Hess's Law: Heats of Reaction

### Problem

Using the following thermochemical data, calculate  $\Delta H^\circ$  for:




<u>Thermochemical Equation</u>	<u><math>\Delta H^\circ</math> (kJ)</u>
1. $\text{Ca}(\text{s}) + 2 \text{C}(\text{graphite}) \rightarrow \text{CaC}_2(\text{s})$	-59.8
2. $\text{Ca}(\text{s}) + \frac{1}{2} \text{O}_2(\text{g}) \rightarrow \text{CaO}(\text{s})$	-635.09
3. $\text{CaO}(\text{s}) + \text{H}_2\text{O}(\text{l}) \rightarrow \text{Ca}(\text{OH})_2(\text{aq})$	-653.1
4. $\text{C}_2\text{H}_2(\text{g}) + \frac{5}{2} \text{O}_2(\text{g}) \rightarrow 2 \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$	-1300.
5. $\text{C}(\text{graphite}) + \text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g})$	-393.509

### Approach





We need to add a series of reactions for which  $\Delta H^\circ$  is known, to yield the overall reaction of interest. We can manipulate the known reactions as we do this. If necessary, reverse some of the known thermochemical

equations so that the major reactants and products of the desired equation are on the appropriate sides. Then, if necessary multiply the known equations by appropriate coefficients so the major reactants and products have the same coefficients as in the desired equation. Add the known equations and the values of  $\Delta H^\circ$  to obtain  $\Delta H^\circ$  for:



### Solution

The first, third, and fourth equations include the major reactants and products, yet equations 1 and 4 must be reversed so that the reactants and products are on the appropriate sides according to the desired equation. Only multipliers are needed, as the coefficients are the same in the known equations as in the desired equation.




This reaction is exothermic and has a  $\Delta H^\circ = -715 \text{ kJ}$ .

## 6. Hess's Law

### Problem

The following data are the thermochemical equations for iron oxides. Find  $\Delta H^\circ$  for:




### Approach

We need to add a series of reactions for which  $\Delta H^\circ$  is known, to yield the overall reaction of interest. We can manipulate the known reactions as we do this. If necessary, reverse some of the known thermochemical equations so that the major reactants and products of the desired equation are on the appropriate sides. Then, if necessary multiply the known equations by appropriate coefficients so the major reactants and products have the same coefficients as in the desired equation. Add the known equations and the values of  $\Delta H^\circ$  to obtain  $\Delta H^\circ$  for:



### Solution

Equation 1 must be reversed and multiplied by 4 so that the reactants and products are on the appropriate side and the coefficients are the same as in the desired equation.



This reaction is exothermic and has  $\Delta H^\circ = -30.4 \text{ kJ}$ .

## 7. Heats of Reaction

### Problem


Given that the standard heat of formation of  $\text{PbO(s)}$  at 298 K is  $-218.99 \text{ kJ/mol}$ , write the thermochemical equation for the formation of  $\text{PbO(s)}$ .

### Approach


The equation must use the elements in their standard states at 298 K as reactants, and one mole of  $\text{PbO(s)}$  in its standard state as the product. In their standard states, lead is a solid and oxygen is gas.

### Solution

The unbalanced reaction is:



Now, balance the equation by putting a coefficient of  $1/2$  before  $\text{O}_2$ . Remember: we must form a single mole of  $\text{PbO}_2$ .



The heat of reaction of  $\text{PbO(s)}$  is given as  $-218.99 \text{ kJ/mol}$ , so the complete reaction is:



## 8. Calorimetry

### Problem

A 40.00 mg sample of naphthalene,  $\text{C}_{10}\text{H}_8$ , was combusted in a calorimeter containing 750. g  $\text{H}_2\text{O}$ . The temperature of the calorimeter increased from  $25.000^\circ\text{C}$  to  $25.404^\circ\text{C}$ . The calorimeter has a heat capacity (calorimeter constant) of  $465 \text{ J/}^\circ\text{C}$ . Calculate  $\Delta H^\circ_{\text{combustion}}$  for naphthalene in  $\text{kJ/mol}$ .

### Approach

First, calculate the energy to heat the water,  $q_{\text{water}}$ . Then, calculate the energy to heat the calorimeter vessel,  $q_{\text{calorimeter}}$ . The total energy transferred is the sum of these two numbers,  $q_{\text{system}}$ . Next, convert the mass of the naphthalene sample into moles. Finally, solve for the heat transfer using  $q_{\text{system}}$  and the moles of naphthalene.

## Solution

**Step 1.** Solve for  $q_{\text{water}}$ .

$$q(\text{J})_{\text{water}} = 750. \text{ g} * 4.184 \text{ J/g K} * 0.404 \text{ K} = 1.27 \times 10^3 \text{ J}$$

**Step 2.** Solve for  $q_{\text{calorimeter}}$ .

$$q(\text{J})_{\text{calorimeter}} = 465 \text{ J/K} * 0.404 \text{ K} = 1.88 \times 10^2 \text{ J}$$

**Step 3.** Solve for  $q_{\text{system}}$ , which is the negative of the total energy transferred.

$$\begin{aligned} \text{Total energy transferred} &= -q(\text{J})_{\text{system}} \\ &= -(1.27 \times 10^3 \text{ J} + 1.88 \times 10^2 \text{ J}) = -1.46 \text{ kJ} \end{aligned}$$

**Step 4.** Convert mass of naphthalene to moles naphthalene using the molar mass.

$$\begin{aligned} 0.0400 \text{ g naphthalene} &* 1 \text{ mol naphthalene} / 128.17 \text{ g naphthalene} \\ &= 3.12 \times 10^{-4} \text{ mol naphthalene} \end{aligned}$$

**Step 5.** Finally, solve for heat transfer by dividing  $q_{\text{system}}$  in kilojoules by the moles of naphthalene.

$$\Delta H^\circ = -1.46 \text{ kJ} / 3.12 \times 10^{-4} \text{ mol naphthalene} = -4.66 \times 10^3 \text{ kJ/mol}$$