

CH 221 Problem Set #5

Complete problem set on separate pieces of paper showing all work, circling final answers, etc.

Covering: Chapter Three (3.3-3.4), Chapter Five and Chapter Guide Five


Important Tables / Constants: $C(\text{H}_2\text{O}) = 4.184 \text{ J g}^{-1} \text{ K}^{-1}$, $\log_{10} x = \ln x / \ln 10$ and the *Thermodynamic Values* found after this problem set and here: <http://mhchem.org/thermo>

- Determine the oxidation number of each element in the following ions or compounds.
 - $\text{O}_2(\text{g})$
 - CuO
 - UO^{2+}
 - $\text{H}_2\text{AsO}_4^{-1}$
 - OF_2
 - XeO_4^{2-}
- Which of the following reactions are oxidation-reduction reactions? Explain your answer briefly. Classify the remaining reactions.
 - $\text{CdCl}_2(\text{aq}) + \text{Na}_2\text{S}(\text{aq}) \rightarrow \text{CdS}(\text{s}) + 2 \text{NaCl}(\text{aq})$
 - $2 \text{Ca}(\text{s}) + \text{O}_2(\text{g}) \rightarrow 2 \text{CaO}(\text{s})$
 - $4 \text{Fe}(\text{OH})_2(\text{aq}) + 2 \text{H}_2\text{O}(\ell) + \text{O}_2(\text{g}) \rightarrow 4 \text{Fe}(\text{OH})_3(\text{aq})$
 - $\text{MgCO}_3(\text{s}) + 2 \text{HCl}(\text{aq}) \rightarrow \text{MgCl}_2(\text{aq}) + \text{H}_2\text{O}(\ell) + \text{CO}_2(\text{g})$
- In the following reactions, decide which reactant is oxidized and which is reduced. Designate the oxidizing agent and the reducing agent.
 - $\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 3 \text{Sn}^{2+}(\text{aq}) + 14 \text{H}^+(\text{aq}) \rightarrow 2 \text{Cr}^{3+}(\text{aq}) + 3 \text{Sn}^{4+}(\text{aq}) + 7 \text{H}_2\text{O}(\ell)$
 - $\text{FeS}(\text{s}) + 3 \text{NO}_3^{-1}(\text{aq}) + 4 \text{H}^+(\text{aq}) \rightarrow 3 \text{NO}(\text{g}) + \text{SO}_4^{2-}(\text{aq}) + \text{Fe}^{3+}(\text{aq}) + 2 \text{H}_2\text{O}(\ell)$
- What is the mass of solute, in grams, in 125 mL of a $1.023 \times 10^{-3} \text{ M}$ solution of Na_3PO_4 ? What are the molar concentrations of Na^+ and PO_4^{3-} ions?
- Twelve (12.0) mL of a 0.125 M BaCl_2 solution is diluted with 9.0 mL of water to a total volume of 21.0 mL. What is the concentration (M) of the new solution? What is the concentration (M) of Cl^{-1} in the final solution? How many grams of BaCl_2 are in the final solution?
- A saturated solution of milk of magnesia, $\text{Mg}(\text{OH})_2$, has a pH of 10.5. What is the hydrogen ion concentration of the solution? Is the solution acidic or basic?
- What mass of Na_2CO_3 , in grams, is required for complete reaction with 50.0 mL of 0.125 M HNO_3 ?
 $\text{Na}_2\text{CO}_3(\text{aq}) + 2 \text{HNO}_3(\text{aq}) \rightarrow 2 \text{NaNO}_3(\text{aq}) + \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\ell)$
- Potassium hydrogen phthalate, $\text{KHC}_8\text{H}_4\text{O}_4$, is used to standardize solutions of bases. The acidic anion reacts with strong bases according to the net ionic equation shown below. If a 0.902 g sample of potassium hydrogen phthalate is dissolved in water and titrated to the equivalence point with 26.45 mL of NaOH , what is the molar concentration of the NaOH ?
 $\text{HC}_8\text{H}_4\text{O}_4^{-1}(\text{aq}) + \text{OH}^{-}(\text{aq}) \rightarrow \text{C}_8\text{H}_4\text{O}_4^{-2}(\text{aq}) + \text{H}_2\text{O}(\ell)$
- What quantity of heat is required to raise the temperature of 50.00 mL of water from 25.52 °C to 28.75 °C? The density of water at this temperature is 0.997 g/mL.
- After absorbing 1.850 kJ of heat, the temperature of a 0.500 kg block of copper is 37 °C. What was the initial temperature of the copper? $C_{\text{Cu}} = 0.385 \text{ J/g}^\circ\text{C}$



Problem Set #5 continues on next page

Problem Set #5, Continued from previous page


11. A 45.5 g sample of copper at 99.8 °C is dropped into a beaker containing 152 g of water at 18.5 °C. What is the final temperature when thermal equilibrium is reached? $C_{\text{Cu}} = 0.385 \text{ J/g}^\circ\text{C}$
12. A piece of chromium metal with a mass of 24.26 g is heated in boiling water to 98.3 °C and then dropped into a coffee cup calorimeter containing 82.3 g of water at 23.3 °C. When thermal equilibrium is reached, the final temperature is 25.6 °C. Calculate the specific heat of the chromium.
13. Chloromethane, CH_3Cl , arises from the oceans and from microbial fermentation and is found throughout the environment. It is used in the manufacture of various chemicals and has been used as a topical anesthetic. What quantity of heat must be absorbed to convert 92.5 g of liquid to a vapor at its boiling point, -24.09°C ? The heat of vaporization of CH_3Cl is 21.40 kJ/mol.
14. Calcium carbide, CaC_2 , is manufactured by the reaction of CaO with carbon at high temperatures. Calcium carbide can then be used to make acetylene. Using the reaction below, is this reaction endothermic or exothermic? If 10.0 g of CaO is allowed to react with an excess of carbon, what quantity of heat is absorbed or evolved by the reaction?




15. The enthalpy changes of the following reactions can be measured:



Use these values and Hess's law to determine the enthalpy change for the reaction:



16. You wish to know the enthalpy change for the formation of liquid PCl_3 from the elements:



The enthalpy change for the formation of solid PCl_5 from the elements can be determined experimentally, as can the enthalpy change for the reaction of $\text{PCl}_3\text{(l)}$ with more chlorine to give $\text{PCl}_5\text{(s)}$. Use the equations below to calculate the enthalpy change for the formation of 1.00 mol $\text{PCl}_3\text{(l)}$ from phosphorus and chlorine.



17. Write a balanced chemical equation for the formation of $\text{CaCO}_3\text{(s)}$ from the elements in their standard states. Find the value of ΔH_f° for $\text{CaCO}_3\text{(s)}$ in the appendix of your textbook. If 10.0 g of $\text{CaCO}_3\text{(s)}$ forms from the elements, how much energy is required or will be released?
18. The first step in the production of nitric acid from ammonia involves the oxidation of NH_3 :
- $$4 \text{NH}_3\text{(g)} + 5 \text{O}_2\text{(g)} \rightarrow 4 \text{NO(g)} + 6 \text{H}_2\text{O(g)}$$
- Use standard enthalpies of formation to calculate the standard enthalpy change for this reaction.
 - Using this reaction, what quantity of heat is evolved or absorbed in the *formation* of 10.0 g of NH_3 ?

Standard Thermodynamic Properties for Selected Substances

Substance	ΔH_f° (kJ/mol)	ΔG_f° (kJ/mol)	S° (J/K·mol)
aluminum			
Al(s)	0	0	28.3
Al(g)	324.4	285.7	164.54
Al ₂ O ₃ (s)	-1676	-1582	50.92
AlF ₃ (s)	-1510.4	-1425	66.5
AlCl ₃ (s)	-704.2	-628.8	110.67
AlCl ₃ ·6H ₂ O(s)	-2691.57	-2269.40	376.56
Al ₂ S ₃ (s)	-724.0	-492.4	116.9
Al ₂ (SO ₄) ₃ (s)	-3445.06	-3506.61	239.32
antimony			
Sb(s)	0	0	45.69
Sb(g)	262.34	222.17	180.16
Sb ₄ O ₆ (s)	-1440.55	-1268.17	220.92
SbCl ₃ (g)	-313.8	-301.2	337.80
SbCl ₅ (g)	-394.34	-334.29	401.94
Sb ₂ S ₃ (s)	-174.89	-173.64	182.00
SbCl ₃ (s)	-382.17	-323.72	184.10
SbOCl(s)	-374.0	-	-
arsenic			
As(s)	0	0	35.1
As(g)	302.5	261.0	174.21
As ₄ (g)	143.9	92.4	314
As ₄ O ₆ (s)	-1313.94	-1152.52	214.22
As ₂ O ₅ (s)	-924.87	-782.41	105.44
AsCl ₃ (g)	-261.50	-248.95	327.06
As ₂ S ₃ (s)	-169.03	-168.62	163.59
AsH ₃ (g)	66.44	68.93	222.78
H ₃ AsO ₄ (s)	-906.3	—	—
barium			
Ba(s)	0	0	62.5
Ba(g)	180	146	170.24
BaO(s)	-548.0	-520.3	72.1
BaCl ₂ (s)	-855.0	-806.7	123.7
BaSO ₄ (s)	-1473.2	-1362.3	132.2
beryllium			
Be(s)	0	0	9.50
Be(g)	324.3	286.6	136.27
BeO(s)	-609.4	-580.1	13.8
bismuth			
Bi(s)	0	0	56.74
Bi(g)	207.1	168.2	187.00
Bi ₂ O ₃ (s)	-573.88	-493.7	151.5
BiCl ₃ (s)	-379.07	-315.06	176.98
Bi ₂ S ₃ (s)	-143.1	-140.6	200.4
boron			
B(s)	0	0	5.86
B(g)	565.0	521.0	153.4
B ₂ O ₃ (s)	-1273.5	-1194.3	53.97
B ₂ H ₆ (g)	36.4	87.6	232.1
H ₃ BO ₃ (s)	-1094.33	-968.92	88.83
BF ₃ (g)	-1136.0	-1119.4	254.4
BCl ₃ (g)	-403.8	-388.7	290.1
B ₃ N ₃ H ₆ (l)	-540.99	-392.79	199.58

Standard Thermodynamic Properties for Selected Substances

Substance	ΔH_f° (kJ/mol)	ΔG_f° (kJ/mol)	S° (J/K·mol)
boron continued			
HBO ₂ (s)	-794.25	-723.41	37.66
bromine			
Br ₂ (l)	0	0	152.23
Br ₂ (g)	30.91	3.142	245.5
Br(g)	111.88	82.429	175.0
BrF ₃ (g)	-255.60	-229.45	292.42
HBr(g)	-36.3	-53.43	198.7
cadmium			
Cd(s)	0	0	51.76
Cd(g)	112.01	77.41	167.75
CdO(s)	-258.2	-228.4	54.8
CdCl ₂ (s)	-391.5	-343.9	115.3
CdSO ₄ (s)	-933.3	-822.7	123.0
CdS(s)	-161.9	-156.5	64.9
calcium			
Ca(s)	0	0	41.6
Ca(g)	178.2	144.3	154.88
CaO(s)	-634.9	-603.3	38.1
Ca(OH) ₂ (s)	-985.2	-897.5	83.4
CaSO ₄ (s)	-1434.5	-1322.0	106.5
CaSO ₄ ·2H ₂ O(s)	-2022.63	-1797.45	194.14
CaCO ₃ (s) (calcite)	-1220.0	-1081.4	110.0
CaSO ₃ ·H ₂ O(s)	-1752.68	-1555.19	184.10
carbon			
C(s) (graphite)	0	0	5.740
C(s) (diamond)	1.89	2.90	2.38
C(g)	716.681	671.2	158.1
CO(g)	-110.52	-137.15	197.7
CO ₂ (g)	-393.51	-394.36	213.8
CH ₄ (g)	-74.6	-50.5	186.3
CH ₃ OH(l)	-239.2	-166.6	126.8
CH ₃ OH(g)	-201.0	-162.3	239.9
CCl ₄ (l)	-128.2	-62.5	214.4
CCl ₄ (g)	-95.7	-58.2	309.7
CHCl ₃ (l)	-134.1	-73.7	201.7
CHCl ₃ (g)	-103.14	-70.34	295.71
CS ₂ (l)	89.70	65.27	151.34
CS ₂ (g)	116.9	66.8	238.0
C ₂ H ₂ (g)	227.4	209.2	200.9
C ₂ H ₄ (g)	52.4	68.4	219.3
C ₂ H ₆ (g)	-84.0	-32.0	229.2
CH ₃ CO ₂ H(l)	-484.3	-389.9	159.8
CH ₃ CO ₂ H(g)	-434.84	-376.69	282.50
C ₂ H ₅ OH(l)	-277.6	-174.8	160.7
C ₂ H ₅ OH(g)	-234.8	-167.9	281.6
C ₃ H ₈ (g)	-103.8	-23.4	270.3
C ₆ H ₆ (g)	82.927	129.66	269.2
C ₆ H ₆ (l)	49.1	124.50	173.4
CH ₂ Cl ₂ (l)	-124.2	-63.2	177.8
CH ₂ Cl ₂ (g)	-95.4	-65.90	270.2
CH ₃ Cl(g)	-81.9	-60.2	234.6

Standard Thermodynamic Properties for Selected Substances

Substance	ΔH_f° (kJ/mol)	ΔG_f° (kJ/mol)	S° (J/K·mol)
carbon continued			
C ₂ H ₅ Cl(l)	-136.52	-59.31	190.79
C ₂ H ₅ Cl(g)	-112.17	-60.39	276.00
C ₂ N ₂ (g)	308.98	297.36	241.90
HCN(l)	108.9	125.0	112.8
HCN(g)	135.5	124.7	201.8
chlorine			
Cl ₂ (g)	0	0	223.1
Cl(g)	121.3	105.70	165.2
ClF(g)	-54.48	-55.94	217.78
ClF ₃ (g)	-158.99	-118.83	281.50
Cl ₂ O(g)	80.3	97.9	266.2
Cl ₂ O ₇ (l)	238.1	—	—
Cl ₂ O ₇ (g)	272.0	—	—
HCl(g)	-92.307	-95.299	186.9
HClO ₄ (l)	-40.58	—	—
chromium			
Cr(s)	0	0	23.77
Cr(g)	396.6	351.8	174.50
Cr ₂ O ₃ (s)	-1139.7	-1058.1	81.2
CrO ₃ (s)	-589.5	—	—
(NH ₄) ₂ Cr ₂ O ₇ (s)	-1806.7	—	—
cobalt			
Co(s)	0	0	30.0
CoO(s)	-237.9	-214.2	52.97
Co ₃ O ₄ (s)	-910.02	-794.98	114.22
Co(NO ₃) ₂ (s)	-420.5	—	—
copper			
Cu(s)	0	0	33.15
Cu(g)	338.32	298.58	166.38
CuO(s)	-157.3	-129.7	42.63
Cu ₂ O(s)	-168.6	-146.0	93.14
CuS(s)	-53.1	-53.6	66.5
Cu ₂ S(s)	-79.5	-86.2	120.9
CuSO ₄ (s)	-771.36	-662.2	109.2
Cu(NO ₃) ₂ (s)	-302.9	—	—
fluorine			
F ₂ (g)	0	0	202.8
F(g)	79.4	62.3	158.8
F ₂ O(g)	24.7	41.9	247.43
HF(g)	-273.3	-275.4	173.8
hydrogen			
H ₂ (g)	0	0	130.7
H(g)	217.97	203.26	114.7
H ₂ O(l)	-285.83	-237.1	70.0
H ₂ O(g)	-241.82	-228.59	188.8
H ₂ O ₂ (l)	-187.78	-120.35	109.6
H ₂ O ₂ (g)	-136.3	-105.6	232.7
HF(g)	-273.3	-275.4	173.8
HCl(g)	-92.307	-95.299	186.9
HBr(g)	-36.3	-53.43	198.7
HI(g)	26.48	1.70	206.59

Standard Thermodynamic Properties for Selected Substances

Substance	ΔH_f° (kJ/mol)	ΔG_f° (kJ/mol)	S° (J/K·mol)
hydrogen continued			
H ₂ S(g)	-20.6	-33.4	205.8
H ₂ Se(g)	29.7	15.9	219.0
iodine			
I ₂ (s)	0	0	116.14
I ₂ (g)	62.438	19.3	260.7
I(g)	106.84	70.2	180.8
IF(g)	95.65	-118.49	236.06
ICl(g)	17.78	-5.44	247.44
IBr(g)	40.84	3.72	258.66
IF ₇ (g)	-943.91	-818.39	346.44
HI(g)	26.48	1.70	206.59
iron			
Fe(s)	0	0	27.3
Fe(g)	416.3	370.7	180.5
Fe ₂ O ₃ (s)	-824.2	-742.2	87.40
Fe ₃ O ₄ (s)	-1118.4	-1015.4	146.4
Fe(CO) ₅ (l)	-774.04	-705.42	338.07
Fe(CO) ₅ (g)	-733.87	-697.26	445.18
FeCl ₂ (s)	-341.79	-302.30	117.95
FeCl ₃ (s)	-399.49	-334.00	142.3
FeO(s)	-272.0	-255.2	60.75
Fe(OH) ₂ (s)	-569.0	-486.5	88.
Fe(OH) ₃ (s)	-823.0	-696.5	106.7
FeS(s)	-100.0	-100.4	60.29
Fe ₃ C(s)	25.10	20.08	104.60
lead			
Pb(s)	0	0	64.81
Pb(g)	195.2	162.	175.4
PbO(s) (yellow)	-217.32	-187.89	68.70
PbO(s) (red)	-218.99	-188.93	66.5
Pb(OH) ₂ (s)	-515.9	—	—
PbS(s)	-100.4	-98.7	91.2
Pb(NO ₃) ₂ (s)	-451.9	—	—
PbO ₂ (s)	-277.4	-217.3	68.6
PbCl ₂ (s)	-359.4	-314.1	136.0
lithium			
Li(s)	0	0	29.1
Li(g)	159.3	126.6	138.8
LiH(s)	-90.5	-68.3	20.0
LiOH(s)	-487.5	-441.5	42.8
LiF(s)	-616.0	-587.5	35.7
Li ₂ CO ₃ (s)	-1216.04	-1132.19	90.17
manganese			
Mn(s)	0	0	32.0
Mn(g)	280.7	238.5	173.7
MnO(s)	-385.2	-362.9	59.71
MnO ₂ (s)	-520.03	-465.1	53.05
Mn ₂ O ₃ (s)	-958.97	-881.15	110.46
Mn ₃ O ₄ (s)	-1378.83	-1283.23	155.64
mercury			
Hg(l)	0	0	75.9

Standard Thermodynamic Properties for Selected Substances

Substance	ΔH_f° (kJ/mol)	ΔG_f° (kJ/mol)	S° (J/K·mol)
mercury continued			
Hg(g)	61.4	31.8	175.0
HgO(s) (red)	-90.83	-58.5	70.29
HgO(s) (yellow)	-90.46	-58.43	71.13
HgCl ₂ (s)	-224.3	-178.6	146.0
Hg ₂ Cl ₂ (s)	-265.4	-210.7	191.6
HgS(s) (red)	-58.16	-50.6	82.4
HgS(s) (black)	-53.56	-47.70	88.28
HgSO ₄ (s)	-707.51	-594.13	0.00
nitrogen			
N ₂ (g)	0	0	191.6
N(g)	472.704	455.5	153.3
NO(g)	90.25	87.6	210.8
NO ₂ (g)	33.2	51.30	240.1
N ₂ O(g)	81.6	103.7	220.0
N ₂ O ₃ (g)	83.72	139.41	312.17
N ₂ O ₄ (g)	11.1	99.8	304.4
N ₂ O ₅ (g)	11.3	115.1	355.7
NH ₃ (g)	-45.9	-16.5	192.8
N ₂ H ₄ (l)	50.63	149.43	121.21
N ₂ H ₄ (g)	95.4	159.4	238.5
NH ₄ NO ₃ (s)	-365.56	-183.87	151.08
NH ₄ Cl(s)	-314.43	-202.87	94.6
NH ₄ Br(s)	-270.8	-175.2	113.0
NH ₄ I(s)	-201.4	-112.5	117.0
NH ₄ NO ₂ (s)	-256.5	—	—
HNO ₃ (l)	-174.1	-80.7	155.6
HNO ₃ (g)	-133.9	-73.5	266.9
oxygen			
O ₂ (g)	0	0	205.2
O(g)	249.17	231.7	161.1
O ₃ (g)	142.7	163.2	238.9
phosphorus			
P ₄ (s)	0	0	164.4
P ₄ (g)	58.91	24.4	280.0
P(g)	314.64	278.25	163.19
PH ₃ (g)	5.4	13.5	210.2
PCl ₃ (g)	-287.0	-267.8	311.78
PCl ₅ (g)	-374.9	-305.0	364.4
P ₄ O ₆ (s)	-1640.1	—	—
P ₄ O ₁₀ (s)	-2984.0	-2697.0	228.86
HPO ₃ (s)	-948.5	—	—
H ₃ PO ₂ (s)	-604.6	—	—
H ₃ PO ₃ (s)	-964.4	—	—
H ₃ PO ₄ (s)	-1279.0	-1119.1	110.50
H ₃ PO ₄ (l)	-1266.9	-1124.3	110.5
H ₄ P ₂ O ₇ (s)	-2241.0	—	—
POCl ₃ (l)	-597.1	-520.8	222.5
POCl ₃ (g)	-558.5	-512.9	325.5
potassium			
K(s)	0	0	64.7
K(g)	89.0	60.5	160.3

Standard Thermodynamic Properties for Selected Substances

Substance	ΔH_f° (kJ/mol)	ΔG_f° (kJ/mol)	S° (J/K·mol)
potassium continued			
KF(s)	-576.27	-537.75	66.57
KCl(s)	-436.5	-408.5	82.6
silicon			
Si(s)	0	0	18.8
Si(g)	450.0	405.5	168.0
SiO ₂ (s)	-910.7	-856.3	41.5
SiH ₄ (g)	34.3	56.9	204.6
H ₂ SiO ₃ (s)	-1188.67	-1092.44	133.89
H ₄ SiO ₄ (s)	-1481.14	-1333.02	192.46
SiF ₄ (g)	-1615.0	-1572.8	282.8
SiCl ₄ (l)	-687.0	-619.8	239.7
SiCl ₄ (g)	-662.75	-622.58	330.62
SiC(s, beta cubic)	-73.22	-70.71	16.61
SiC(s, alpha hexagonal)	-71.55	-69.04	16.48
silver			
Ag(s)	0	0	42.55
Ag(g)	284.9	246.0	172.89
Ag ₂ O(s)	-31.05	-11.20	121.3
AgCl(s)	-127.0	-109.8	96.3
Ag ₂ S(s)	-32.6	-40.7	144.0
sodium			
Na(s)	0	0	51.3
Na(g)	107.5	77.0	153.7
Na ₂ O(s)	-414.2	-375.5	75.1
NaCl(s)	-411.2	-384.1	72.1
sulfur			
S ₈ (s) (rhombic)	0	0	256.8
S(g)	278.81	238.25	167.82
SO ₂ (g)	-296.83	-300.1	248.2
SO ₃ (g)	-395.72	-371.06	256.76
H ₂ S(g)	-20.6	-33.4	205.8
H ₂ SO ₄ (l)	-813.989	690.00	156.90
H ₂ S ₂ O ₇ (s)	-1273.6	—	—
SF ₄ (g)	-728.43	-684.84	291.12
SF ₆ (g)	-1220.5	-1116.5	291.5
SCl ₂ (l)	-50	—	—
SCl ₂ (g)	-19.7	—	—
S ₂ Cl ₂ (l)	-59.4	—	—
S ₂ Cl ₂ (g)	-19.50	-29.25	319.45
SOCl ₂ (g)	-212.55	-198.32	309.66
SOCl ₂ (l)	-245.6	—	—
SO ₂ Cl ₂ (l)	-394.1	—	—
SO ₂ Cl ₂ (g)	-354.80	-310.45	311.83
tin			
Sn(s)	0	0	51.2
Sn(g)	301.2	266.2	168.5
SnO(s)	-285.8	-256.9	56.5
SnO ₂ (s)	-577.6	-515.8	49.0
SnCl ₄ (l)	-511.3	-440.1	258.6
SnCl ₄ (g)	-471.5	-432.2	365.8

Standard Thermodynamic Properties for Selected Substances

Substance	ΔH_f° (kJ/mol)	ΔG_f° (kJ/mol)	S° (J/K·mol)
titanium			
Ti(s)	0	0	30.7
Ti(g)	473.0	428.4	180.3
TiO ₂ (s)	-944.0	-888.8	50.6
TiCl ₄ (l)	-804.2	-737.2	252.4
TiCl ₄ (g)	-763.2	-726.3	353.2
tungsten			
W(s)	0	0	32.6
W(g)	849.4	807.1	174.0
WO ₃ (s)	-842.9	-764.0	75.9
zinc			
Zn(s)	0	0	41.6
Zn(g)	130.73	95.14	160.98
ZnO(s)	-350.5	-320.5	43.7
ZnCl ₂ (s)	-415.1	-369.43	111.5
ZnS(s)	-206.0	-201.3	57.7
ZnSO ₄ (s)	-982.8	-871.5	110.5
ZnCO ₃ (s)	-812.78	-731.57	82.42
complexes			
<i>cis</i> -[Co(NH ₃) ₄ (NO ₂) ₂]NO ₃	-898.7	—	—
<i>trans</i> -[Co(NH ₃) ₄ (NO ₂) ₂]NO ₃	-896.2	—	—
NH ₄ [Co(NH ₃) ₂ (NO ₂) ₄]	-837.6	—	—
[Co(NH ₃) ₆][Co(NH ₃) ₂ (NO ₂) ₄] ₃	-2733.0	—	—
<i>cis</i> -[Co(NH ₃) ₄ Cl ₂]Cl	-874.9	—	—
<i>trans</i> -[Co(NH ₃) ₄ Cl ₂]Cl	-877.4	—	—
<i>cis</i> -[Co(en) ₂ (NO ₂) ₂]NO ₃	-689.5	—	—
<i>cis</i> -[Co(en) ₂ Cl ₂]Cl	-681.2	—	—
<i>trans</i> -[Co(en) ₂ Cl ₂]Cl	-677.4	—	—
[Co(en) ₃](ClO ₄) ₃	-762.7	—	—
[Co(en) ₃]Br ₂	-595.8	—	—
[Co(en) ₃]I ₂	-475.3	—	—
[Co(en) ₃]I ₃	-519.2	—	—
[Co(NH ₃) ₆](ClO ₄) ₃	-1034.7	-221.1	615
[Co(NH ₃) ₅ NO ₂](NO ₃) ₂	-1088.7	-412.9	331
[Co(NH ₃) ₆](NO ₃) ₃	-1282.0	-524.5	448
[Co(NH ₃) ₅ Cl]Cl ₂	-1017.1	-582.5	366.1
[Pt(NH ₃) ₄]Cl ₂	-725.5	—	—
[Ni(NH ₃) ₆]Cl ₂	-994.1	—	—
[Ni(NH ₃) ₆]Br ₂	-923.8	—	—
[Ni(NH ₃) ₆]I ₂	-808.3	—	—

Standard Thermodynamic Properties for Selected Substances

Blank page for printing purposes!