

Chemistry 221 Exam II Review Chapters 3, 4 and 5



Chemistry 221
Professor Michael Russell



Which statement regarding this reaction is true?

- A. 3 moles of S are produced per mole of H_2S .
- B. 1 mole of SO_2 is consumed per mole of H_2S .
- C. 1 mole of H_2O is produced per mole of H_2S .
- D. The total number of moles of products is always equal to the total number of moles of reactants used.
- E. None of these statements are true.

What is the balanced equation for the combustion of butane, C_4H_{10} ?

- A. $\text{C}_4\text{H}_{10}\text{(g)} + \text{O}_2\text{(g)} \rightarrow \text{CO}_2\text{(g)} + \text{H}_2\text{O(g)}$
- B. $2 \text{C}_4\text{H}_{10}\text{(g)} + 13 \text{O}_2\text{(g)} \rightarrow 8 \text{CO}_2\text{(g)} + 10 \text{H}_2\text{O(g)}$
- C. $\text{C}_4\text{H}_{10}\text{(g)} + 13 \text{O}_2\text{(g)} \rightarrow 4 \text{CO}_2\text{(g)} + 5 \text{H}_2\text{O(g)}$
- D. $\text{C}_4\text{H}_{10}\text{(g)} + 9 \text{O}_2\text{(g)} \rightarrow 4 \text{CO}_2\text{(g)} + 10 \text{H}_2\text{O(g)}$

In the reaction of 2.0 mol of CCl_4 with an excess of HF, 1.7 mol of CCl_2F_2 is obtained.



Which statement is true here?

- A. The theoretical yield for CCl_2F_2 is 1.7 mol.
- B. The actual yield for CCl_2F_2 is 1.0 mol.
- C. The percent yield for the reaction is 85%.
- D. Theoretical yield cannot be determined unless the exact amount of HF used is known.
- E. Infinite diversity in infinite combinations (IDIC)


Burning sulfur in an atmosphere of fluorine produces the very stable compound SF_6 .



If you wish to produce 2.50 moles of SF_6 , you will need to use:

- A. 0.313 moles of S_8 and 7.50 moles of F_2 .
- B. 1.00 moles of S_8 and 24.0 moles of F_2 .
- C. 0.125 moles of S_8 and 3.00 moles of F_2 .
- D. 8.00 moles of S_8 and 24.0 moles of F_2 .
- E. More information is required to answer this question.

Ammonia is prepared by the reaction:



If 10.0 mol of N_2 are mixed with 25.0 mol of H_2 , the amount of NH_3 produced will be:

- A. 20.0 mol NH_3
- B. 16.7 mol NH_3
- C. 37.5 mol NH_3
- D. 25.0 mol NH_3
- E. 35.0 mol NH_3

A compound with C, H and O is found through combustion analysis of a 0.255 g sample to give 0.561 g CO₂ and 0.306 g H₂O; it also has a molar mass of 60.1 g/mol. What is the molecular formula?

- A. CH₃CO₂H
- B. C₄H₉O₃
- C. C₃H₆O
- D. C₉H₇O₃
- E. C₃H₈O

MAR

Which of the following is the only insoluble salt in water?

- A. NH₄NO₃
- B. NaOH
- C. PbI₂
- D. K₂CO₃
- E. LiCl

MAR

Which of the compounds below is *not* an acid in aqueous solution?

- A. CH₃CO₂H
- B. H₃PO₄
- C. NH₃
- D. HCl
- E. HClO₄

MAR

Which equation below best represents the balanced, net ionic equation for the reaction of magnesium carbonate with nitric acid?

- A. MgCO₃(s) + 2 HNO₃(aq) → Mg(NO₃)₂(aq) + CO₂(g) + H₂O(l)
- B. MgCO₃(s) + 2 H⁺(aq) → Mg²⁺(aq) + CO₂(g) + H₂O(l)
- C. Mg²⁺(aq) + 2 NO₃⁻(aq) → Mg(NO₃)₂(s)
- D. MgCO₃(s) + 2 HNO₃(aq) → Mg(NO₃)₂(aq) + H₂CO₃(aq)
- E. More information is required to answer this question.

MAR

Which equation below best represents the balanced net ionic equation for the reaction of potassium hydroxide and iron(II) chloride to give iron(II) hydroxide and potassium chloride?

- A. 2 KOH(aq) + FeCl₂(aq) → Fe(OH)₂(s) + 2 KCl(aq)
- B. 2 KOH(aq) + FeCl₂(aq) → Fe(OH)₂(aq) + 2 KCl(aq)
- C. 2 OH⁻(aq) + Fe²⁺(aq) → Fe(OH)₂(s)
- D. K⁺(aq) + Cl⁻(aq) → KCl(aq)
- E. More information is required to answer this question.

MAR

Which of the following statements is correct regarding the reaction of Zn with VO₂⁺?



- A. Zn is oxidized and VO₂⁺ is the reducing agent.
- B. Zn is reduced and VO₂⁺ is the reducing agent.
- C. Zn is oxidized and VO₂⁺ is the oxidizing agent.
- D. Zn is reduced and VO₂⁺ is the oxidizing agent.
- E. This is not a redox reaction.

MAR

Assume you dissolve 6.73 g Na_2CO_3 in enough water to make 250. mL of solution. (Molar mass of $\text{Na}_2\text{CO}_3 = 106 \text{ g/mol}$.) What is the concentration of the sodium carbonate?

- A. 26.9 M
- B. 0.0635 M
- C. 0.254 M
- D. 0.762 M
- E. 42 M

MAR

60.0 mL of 0.25 M HCl are added to a 500. mL volumetric flask; water is added to the mark on the flask. What is the concentration of HCl in the diluted solution?

- A. 0.015 M
- B. 0.025 M
- C. 0.030 M
- D. 0.060 M
- E. 0.050 M

MAR

What is the pH of dilute nitric acid with a concentration of 0.030 M?

- A. 0.030
- B. 1.52
- C. 1.82
- D. 2.50
- E. 3.00

MAR

What mass of Na_2CO_3 (molar mass = 106.0 g/mol) is required for complete reaction with 25.0 mL of 0.155 M HNO_3 ?



- A. 0.410 g
- B. 205 g
- C. 0.205 g
- D. 0.122 g
- E. 37 kg

MAR

A piece of copper (5.00 g) is heated for 2.0 seconds, and 100. J of heat energy is transferred to the copper. The temperature increases from 20.0 °C to 71.9 °C. Calculate the specific heat capacity of copper.

- A. 0.278 J/g•K
- B. 0.385 J/g•K
- C. 1.93 J/g•K
- D. 2.60 J/g•K
- E. -0.977 J/g•K

MAR

When 108 grams of water at 22.5 °C are mixed with 65.1 grams of water at an unknown temperature, the final temperature of the mixture is 47.9 °C. What was the initial temperature of the other sample of water?

- A. 8.9 °C
- B. 79.7 °C
- C. 67.0 °C
- D. 90.0 °C
- E. 274 °C

MAR

The standard molar enthalpy of combustion for propane is -2044 kilojoules.



What is the standard enthalpy change for the combustion of 3.000 mol of propane (C_3H_8)?

- A. -6132 kJ
- B. -2044 kJ
- C. -4088 kJ
- D. +2044 kJ
- E. +6132 kJ

MAR

Calculate the enthalpy for the reaction
 $\text{SiH}_4(\text{g}) + 2 \text{O}_2(\text{g}) \rightarrow \text{SiO}_2(\text{g}) + 2 \text{H}_2\text{O}(\text{g})$

using these values:

$$\Delta H_f^\circ[\text{SiH}_4(\text{g})] = +34.3 \text{ kJ/mol};$$

$$\Delta H_f^\circ[\text{SiO}_2(\text{g})] = -910.9 \text{ kJ/mol}; \text{ and}$$

$$\Delta H_f^\circ[\text{H}_2\text{O}(\text{g})] = -241.8 \text{ kJ/mol}$$

- A. -1187.0 kJ/rxn
- B. -1428.8 kJ/rxn
- C. -1360.2 kJ/rxn
- D. -2218.7 kJ/rxn
- E. Not enough information


MAR

Which equation below defines the standard molar enthalpy of formation of gaseous methanol, CH_3OH ?

- A. $\text{CH}_4(\text{g}) + \frac{1}{2} \text{O}_2(\text{g}) \rightarrow \text{CH}_3\text{OH}(\text{g})$
- B. $\text{C}(\text{s}) + 2 \text{H}_2(\text{g}) + \frac{1}{2} \text{O}_2(\text{g}) \rightarrow \text{CH}_3\text{OH}(\text{g})$
- C. $\text{CO}(\text{g}) + 2 \text{H}_2(\text{g}) \rightarrow \text{CH}_3\text{OH}(\text{g})$
- D. $\text{H}_2\text{O}(\text{g}) + \text{C}(\text{s}) + \text{H}_2(\text{g}) \rightarrow \text{CH}_3\text{OH}(\text{g})$
- E. You'll go blind if you drink methanol! Who cares! :)

MAR

Calculate the standard molar enthalpy of formation for $\text{FeCl}_2(\text{s})$ using the following:



- A. -57.7 kJ/mol
- B. -341.8 kJ/mol
- C. -284.1 kJ/mol
- D. -457.2 kJ/mol
- E. 42 kJ/mol

MAR


**End of
Review -
good luck
with your
studying!**



Need more practice?

- Practice Problem Sets (online)
- Concept Guides (Companion and online)
- Chapter Guides (online)
- End of Chapter Problems in Textbook (every other question has answer at end)

Good luck with your studying!



MAR