Chapter 7 Part I: Stoichiometry of Chemical Reactions (7.1 - 7.2 *only*)

Chemistry 221 Professor Michael Russell http://mhchem.org/221 MAR

7.1 - Writing and Balancing Chemical Equations

A balanced chemical equation (or reaction) uses symbolism to represent the identities and the relative quantities of substances undergoing a chemical (or physical) change.

Chemical equations depict the kind of reactants (left) and products (right) and their relative amounts in a reaction.

 $4 \text{ Al(s)} + 3 O_2(g) \rightarrow 2 \text{ Al}_2O_3(s)$


The numbers in the front are called stoichiometric coefficients

The letters (s), (g), (aq) and (l) are the physical states of compounds (solid, gas, liquid and 'dissolved in water' (aq))

MAR


MAR

Reaction of Phosphorus with Cl₂



Notice the stoichiometric coefficients and the physical states of the reactants and products.

Reaction of Iron with Cl2




Evidence of a chemical reaction: heat change, precipitate formation, gas evolution, color change

Reaction of Methane (CH₄) with O₂

MAR

MAR



Use smallest coefficients possible, i.e. 1:2:1:2 (above), not 2:4:2:4, etc.

Chemical Equations

Because the same atoms are present in a reaction at the beginning and at the end, the amount of matter in a system does not change.

This is the Law of the

Conservation of Matter Also known as the Law of Mass Action



Lavoisier

Because of the principle of the conservation of matter,

an equation must be balanced.

It must have the same number of atoms of the same kind on both sides.

MAR


Lavoisier, 1788



Balancing Equations



2 Al(s) + 3 Br₂(liq) ---> Al₂Br₆(s)



Balancing Equations

 $C_3H_8(g) + C_2(g) \rightarrow CO_2(g) + H_2O(g)$

 $B_4H_{10}(g) + O_2(g) \rightarrow B_2O_3(g) + H_2O(g)$

MAR

Balancing Equations - hints

A reaction which is **heated** may include an uppercase Greek letter delta (Δ) over the arrow:

$$\Delta$$
 CaCO₃(s) \rightarrow CaO(s) + CO₂(g)

It is sometimes convenient to use fractions instead of integers as intermediate coefficients.

$$C_2H_6 + \frac{7}{2}O_2 \rightarrow 3 H_2O + 2 CO_2$$
 (balanced)

Generally fractions are not used, so when balance is achieved, all the equation's coefficients are multiplied by a whole number to convert to whole numbers.

 $2 C_2H_6 + 7 O_2 \rightarrow 6 H_2O + 4 CO_2$ (balanced)

MAR

Balancing Equations - More Hints

Balance those atoms which occur in only one compound on each side last (i.e. O_2 in previous examples)

Balance the remaining atoms first. Try not to use fractions as coefficients.

Never change the subscripts on a formula, only change the coefficients

Reduce coefficients to smallest whole integers

Check your answer if uncertain

Check that charges are balanced

MAR Practice, practice, practice!

Aqueous Compounds

Consider: $CaCl_2(aq) + 2 AgNO_3(aq) \rightarrow Ca(NO_3)_2(aq) + 2 AgCl(s)$


When ionic compounds dissolve in water, they dissociate into their constituent ions. *Examples:*

 $CaCl_2(aq) \longrightarrow Ca^{2+}(aq) + 2 Cl^{1-}(aq)$

 $2 \text{ AgNO}_3(aq) \rightarrow 2 \text{ Ag}^{1+}(aq) + 2 \text{ NO}_3^{1-}(aq)$

 $Ca(NO_3)_2(aq) \rightarrow Ca^{2+}(aq) + 2 NO_3^{1-}(aq)$

...but AgCl does not dissolve in water (it stays a solid).



Aqueous Compounds

Consider: $CaCl_2(aq) + 2 AgNO_3(aq) \rightarrow Ca(NO_3)_2(aq) + 2 AgCl(s)$

We can re-write the equation as:

$$Ca^{2+}(aq) + 2 Cl^{1-}(aq) + 2 Ag^{1+}(aq) + 2 NO_3^{1-}(aq) \rightarrow Ca^{2+}(aq) + 2 NO_3^{1-}(aq) + 2 AgCl(s)$$

Notice that $Ca^{2+}(aq)$ and 2 $NO_3^{1-}(aq)$ appear on both sides - they are spectator ions and can be removed from the reaction to get the **net ionic reaction**:

 $2 \; Cl^{1\text{-}}(aq) \; + \; 2 \; Ag^{1\text{+}}(aq) \; \longrightarrow \; 2 \; AgCl(s) \quad \text{or better yet:}$

 $Cl^{1-}(aq) + Ag^{1+}(aq) \rightarrow AgCl(s)$ This is the **net ionic reaction**

MAR No spectator ions in net ionic reactions

Net Ionic Equations



The two CI⁻ ions are SPECTATOR IONS - they do not participate. Could have used NO₃⁻ (via HNO₃)

 $Mg(s) + 2 HCl(aq) \rightarrow H_2(g) + MgCl_2(aq)$

Break up (aq), leave gases, solids and liquids alone, to write: $Mg(s) + 2 H^{+}(aq) + 2 Cl^{-}(aq) \rightarrow H_{2}(g) + Mg^{2+}(aq) + 2 Cl^{-}(aq)$

Cl is a spectator - leave out - write the Net Ionic Equation:

 $Mg(s) + 2 H^{+}(aq) \rightarrow H_{2}(g) + Mg^{2+}(aq)$

MAR

Net Ionic Equations

 $K_2CrO_4(aq) + Pb(NO_3)_2(aq) \rightarrow PbCrO_4(s) + 2 KNO_3(aq)$

Net Ionic Equation:

MAR

MAR

 $Pb^{2+}(aq) + CrO_4^{2-}(aq) \rightarrow PbCrO_4(s)$

K+ and NO3- are spectators

See Net Ionic Reactions Handout



7.2 - Classifying Chemical Reactions - Precipitation

A precipitation reaction is a reaction where dissolved substances create solid products (a precipitate.)
Chemists use a solubility table to know common combinations of chemicals which result in soluble (no solids) or insoluble (solids form) products.

Our solubility table is slightly different than the textbook's version, but similar.

Textbook's Solubility


Guide

No. 1

Organization of 100' organization organization of 100' organization organization

MAR

Water Solubility of Ionic Compounds



Water Solubility Of Ionic Compounds

Many ionic compounds dissolve in water (soluble), but many are insoluble and do not dissolve.

Many ions make compounds soluble all of the time:

Examples: Na+, K+, Li+, NH₄+, NO₃-, ClO₃-, ClO₄-, CH₃CO₂-, and most SO_4^{2-} , Cl-, Br and l-compounds.

Solubility guide:

http://mhchem.org/sol/

Page III-7a-3 / Chapter Seven Part I Lecture Notes

MAR

lonic Compounds in Aqueous Solution

Soluble ionic compounds dissolve in water to make aqueous solutions.

Insoluble ionic compounds stay undissolved in water.



KMnO₄ in water






 $K^+(aq) + MnO_4^-(aq)$

Chemical Reactions In Water

Many reaction types are

Exchange Reactions



The anions exchange places between cations



 $Pb(NO_3)_2(aq) + 2 KI(aq)$ $\rightarrow PbI_2(s) + 2 KNO_3(aq)$

Exchange reactions often called Double Displacement Reactions

Precipitation Reactions

A precipitation reaction is an exchange reaction where a precipitate (i.e. solid) is formed as a product

The "driving force" is the formation of an insoluble compound - a precipitate.

Fe(NO₃)₃(aq) + 3 NaOH(aq) \rightarrow 3 NaNO₃(aq) + Fe(OH)₃(s)



Net ionic equation

 $Fe^{3+}(aq) + 3 OH \cdot (aq) \rightarrow Fe(OH)_3(s)$

MAR

MAR

Acid-Base Reactions

An acid-base reaction is an exchange reaction where an acid and a base create water and a 'salt'

Acids react readily with bases. The "driving force" is the formation of water.

NaOH(aq) + HCl(aq) →

NaCl(aq) + H₂O(liq)

Net ionic equation: $OH(aq) + H(aq) \rightarrow H_2O(liq)$

This applies to ALL reactions of STRONG acids and

Acid-base reactions often called "neutralizations", water and "salt" created

MAR



What is an Acid?

Hydronium often written as just H+


An ${\it acid}$ is a substance which dissolves in water to create the ${\it hydronium\ ion},\ {\it H}_3{\it O}^+.$

Acids that give the maximum amount of hydronium possible are called **strong acids**.

There are five strong acids (important):

HCI hydrochloric acid
HBr hydrobromic acid
HI hydroiodic acid
HNO₃ nitric acid
HCIO₄ perchloric acid





Weak Acids

Most acids do not create the maximum amount of hydronium possible; they are called **weak acids**. Most foods are weak acids!

Assume an acid is weak unless you know it is strong



acetic acid

Examples:

 ${\rm CH_3CO_2H}$ acetic acid ${\rm H_2CO_3}$ carbonic acid ${\rm H_3PO_4}$ phosphoric acid HF hydrofluoric acid

What is a Base?

A base is a metal hydroxide. Hydroxide ions react with hydronium ions to create water (and energy!) Bases that give the maximum amount of hydroxide possible are called strong bases.



There are three strong bases (important):

NaOH	sodium hydroxide
КОН	potassium hydroxide
LiOH	lithium hydroxide

NaOH	NaOH
NaOH	NaOH
NaOH	NαOH

Weak Bases

Most bases do not create the maximum amount of hydroxide possible; they are called weak bases. Most household cleaners are weak bases Assume a base is weak unless you know it is strong



ammonium hydroxide in



Examples:

NH₂	ammon	ııa

Na₂CO₃ sodium carbonate Na₃PO₄ sodium phosphate

 N_2H_4 hydrazine

MAR


MAR

MAR

	Common Acids and	Bases	
Strong Ac	ids (Strong Electrolytes)	Strong	Bases (Strong Electrolytes)
HCI	Hydrochloric acid	LiOH	Lithium hydroxide
HBr	Hydrobromic acid	NaOH	Sodium hydroxide
HI	Hydroiodic acid	KOH	Potassium hydroxide
HNO ₃	Nitric acid		
HClO ₄	Perchloric acid		
Weak Aci	ds (Weak Electrolytes)*	Weak	Base (Weak Electrolyte)
H₃PO₄	Phosphoric acid	NH₃	Ammonia
H_2CO_3	Carbonic acid		
CH₃CO₂H	Acetic acid		17 4 la a 4 u a
H ₂ C ₂ O ₄	Oxalic acid		Know the stron
C ₄ H ₆ O ₆	Tartaric acid		acids & bases
C ₆ H ₈ O ₇	Citric acid		acius & bases

Gas-Forming Reactions

A gas forming reaction is an exchange reaction where one of the products is a gas. Gases often result from unstable products that break down into a gas immediately after the chemicals are mixed. Example:



Carbonic acid (H₂CO₃) is unstable and forms CO₂ & H₂O

 $H_2CO_3(aq) \rightarrow CO_2(g) + H_2O(l)$

so we should write:

MAR

 $CaCO_3(s) + 2 HCl(aq) \rightarrow CaCl_2(aq) + CO_2(g) + H_2O(l)$

Another gas forming species: ammonium hydroxide

 $NH_4OH(aq) \rightarrow NH_3(q) + H_2O(l)$

Be on the lookout for H_2CO_3 and NH₄OH as products! They will break down!

Combustion Reactions

A special example of a gas-forming reaction; not an exchange reaction. Used in quantitative chemistry; high temperatures

Reactants: (often) oxygen (O2) and "something organic" (C, H, sometimes O or N)

Products: water and carbon dioxide (also NO₂ if N present)


Examples:

$$C_2H_4(g) + 3 O_2(g) \rightarrow 2 H_2O(g) + 2 CO_2(g)$$

 $4 C_6H_5NO_2(g) + 29 O_2(g) \rightarrow 10 H_2O(g) + 24 CO_2(g) + 4 NO_2(g)$

Oxidation-Reduction Reactions

An oxidation-reduction reaction (often called a redox reaction) is a reaction where electrons are transferred from one reactant (the reducing agent) to another reactant (the oxidizing agent). Knowing where electrons come from (and go to) important! Very powerful and common reaction type (batteries, making chemicals, breathing!)



 $2 H_2(g) + O_2(g) \rightarrow 2 H_2O(liq)$ MAR





LEO says GER

MAR

LEO says GER

Lose Gain **Electrons** Electrons Oxidized Reduced

 $Zn(s) \rightarrow Zn^{2+} + 2e^{-}$ $Cu^{2+} + 2e^{-} \rightarrow Cu(s)$

Oxidized (Zn is the reducing agent) **Reduced** (Cu²⁺ is the *oxidizing agent*)

Can also use "OIL RIG": OIL = "Oxidation is Losing" (electrons)
RIG = "Reduction is Gaining" (electrons)

MAR

Redox Reactions

Lose Electrons Oxidized Gain Electrons Reduced



MAR



Oxidized species become more positive (lose electrons), reduced species become more negative (gain electrons)

 $Cu^{2+}(aq) + 2 Ag(s)$

In all reactions: if something has

also been reduced:

Cu(s) + 2 Ag+(aq) \rightarrow


been oxidized then something has

Oxidation numbers help visualize electron transfer pathways

Oxidation Numbers

Use oxidation number rules to determine redox activity:

• Atoms in free element have ox. no. = 0 Zn(s), $O_2(g)$, $Br_2(liq)$



- In simple ions, ox. no. = charge on ion -1 for Cl-, +2 for Mg2+
- In compounds, F is always -1, O is -2 (except peroxides (O = -1) and with F) and H is +1 (except hydrides (H = -1))
- Sum of oxidation numbers = 0 for a compound or equals the overall charge for an ion

MAR

Oxidation Numbers

Example: Find the oxidation number for each element in H₂S.

The oxidation number for **H** is +1, H₂S is neutral, so:

Charge on $H_2S = 0 = (2 \times 1) + 1x$

x = -2 the oxidation number for S is -2!

Example: Find the oxidation number for each element in sulfite, SO₃2-

The oxidation number for O is -2, sulfite has a -2 charge, so:

 $-2 = 1x + (3 \times -2)$

x = +4 the oxidation number for S is +4!

 $H - \ddot{S} - H$

Oxidation Numbers

Determining oxidation numbers takes practice

: 0 - cl - 0:

HF H: +1 F: -1

CIO₄-CI: +7 O: -2

MAR

Single Replacement Reactions

Single Replacement reactions are always redox reactions



Examples:

 $Cu(s) + 2 AgNO_3(aq) \rightarrow Cu(NO_3)_2(aq) + 2Ag(s)$ $Mg(s) + 2 HCl(aq) \rightarrow MgCl_2(aq) + H_2(g)$

MAR

Redox Reaction Examples



NO = reducing agent O_2 = oxidizing agent $2 \text{ NO} + \text{O}_2 \rightarrow 2 \text{ NO}_2$



Fe = reducing agent Cl₂ = oxidizing agent 2 Fe + 3 Cl₂ → 2 FeCl₃

reducing agent = oxidized oxidizing agent = reduced

End of Chapter 7 Part 1

See also:

- Chapter Seven Part 1 Study Guide
- Chapter Seven Part 1 Concept Guide
- Important Equations (following this slide)
- End of Chapter Problems (following this slide)





Important Equations, Constants, and Handouts from this Chapter:

- · Know how the solubility guide works
- Know what makes an acid acidic (and bases basic) and strong or weak; know how to use the pH scale
- Know how to write and determine net ionic equations and find spectator ions
- · Know how to use molarity with solution stoichiometry problems

MAR

solubility (use the Net Ionics precipitation, types of

solubility table), reactions, molarity (M)

Solutions: Solute,

solvent, aqueous,

Balancing Equations: Reactants, Products, states of matter (s, I, g, aq), stoichiometric coefficients, Law of Conservation of Matter ("mass action")

Know the five types of reactions: precipitation, acid-base, gas forming, combustion and redox. Know how to determine if something has been oxidized or reduced (and the oxidizing agent and reducing agent)

End of Chapter Problems: Test Yourself

- Predict whether these compounds would be labeled as insoluble or soluble: HCI, NaCI, AgCI
 Predict the products of this precipitation reaction and write the net ionic equation: NiCI₂(aq) + (NH₄)₂S(aq) → ? List any spectator ions.
 In the following reaction, decide which reactant is oxidized and which is the product of the prod
- reduced. Designate the oxidizing agent and the reducing agent. Si(s) + 2 $\text{Cl}_2(g) \rightarrow \text{SiCl}_4(l)$ Identify the ions and their concentration that exist in this aqueous solution:
- 0.25 M (NH₄)₂SO₄

End of Chapter Problems: Answers

- Soluble: HCI(aq), NaCI(aq). Insoluble: AgCI(s)
- NiCl₂(aq) + (NH₂)₂S(aq) → NiS(s) + 2 NH₄Cl(aq) Ni²⁺(aq) + S²-(aq) → NiS(s) Spectator ions: NH₄+¹ and Cl¹
 Si is oxidized and is the reducing agent; Cl₂ is reduced and is the
- oxidizing agent
 4. 0.50 M NH₄+1; 0.25 M SO₄²-

MAR