


### Chemical Reactions Chapter 4 Part 1





Reactants:  $Zn + I_2$  → Product:  $ZnI_2$

Chemistry 221  
Professor Michael Russell

MAR

### Chemistry as Cooking! - the Chemical Reaction


"Recipe" and technique leads to successful creations  
Must know amounts to add, how much will be produced  
Haphazard additions can be disastrous!



Last update:  
4/10/23

### Chemical Equations

Depict the kind of reactants and products and their relative amounts in a reaction.




The numbers in the front are called stoichiometric coefficients

The letters (s), (g), (aq) and (l) are the physical states of compounds.

MAR


### Reaction of Phosphorus with $Cl_2$



Notice the stoichiometric coefficients and the physical states of the reactants and products.

MAR

### Reaction of Iron with $Cl_2$




Evidence of a chemical reaction:

heat change, precipitate formation, gas evolution, color change

MAR

### Chemical Equations



This equation means:

4 Al atoms + 3  $O_2$  molecules  
---give--->  
2 molecules of  $Al_2O_3$

or

4 moles of Al + 3 moles of  $O_2$   
---give--->  
2 moles of  $Al_2O_3$

MAR




## Chemical Equations

Because the same atoms are present in a reaction at the beginning and at the end, the amount of matter in a system does not change.

The Law of the Conservation of Matter

Also known as the Law of Mass Action



MAR

## Chemical Equations / Lavoisier

Because of the principle of the conservation of matter, an equation must be balanced.

It must have the same number of atoms of the same kind on both sides.





Lavoisier, 1788


MAR





## Balancing Equations



MAR



## Balancing Equations



MAR

## Balancing Equations - Hints

Balance those atoms which occur in only one compound on each side last (i.e.  $\text{O}_2$  in previous examples)

Balance the remaining atoms first

Reduce coefficients to smallest whole integers

Check your answer if uncertain

**Helpful but optional:** Check that charges are balanced

MAR

## STOICHIOMETRY

Stoichiometry is the study of the quantitative aspects of chemical reactions.

Stoichiometry rests on the principle of the conservation of matter.



MAR

## Stoichiometry

The balanced chemical equation



implies *all* of the following ratios:

$$\frac{4 \text{ mol Al}}{3 \text{ mol O}_2} \quad \frac{4 \text{ mol Al}}{2 \text{ mol Al}_2\text{O}_3} \quad \frac{3 \text{ mol O}_2}{2 \text{ mol Al}_2\text{O}_3}$$


$$\frac{3 \text{ mol O}_2}{4 \text{ mol Al}} \quad \frac{2 \text{ mol Al}_2\text{O}_3}{4 \text{ mol Al}} \quad \frac{2 \text{ mol Al}_2\text{O}_3}{3 \text{ mol O}_2}$$

*These are nothing more than "conversion units" in dimensional analysis!*

MAR



MAR

**PROBLEM:** If 454 g of  $\text{NH}_4\text{NO}_3$  decomposes, how much  $\text{N}_2\text{O}$  and  $\text{H}_2\text{O}$  are formed? What is the theoretical yield of products?



**STEP 1**

Write the balanced chemical equation




**STEP 2** Convert mass reactant (454 g)  $\rightarrow$  moles

$$454 \text{ g} \cdot \frac{1 \text{ mol}}{80.04 \text{ g}} = 5.68 \text{ mol NH}_4\text{NO}_3$$

*80.04 g/mol = molar mass of  $\text{NH}_4\text{NO}_3$*


MAR

MAR




**STEP 3** Convert moles reactant  $\rightarrow$  moles product

Relate moles  $\text{NH}_4\text{NO}_3$  to moles product expected.



Express as a **STOICHIOMETRIC FACTOR**:

$$\frac{2 \text{ mol H}_2\text{O produced}}{1 \text{ mol NH}_4\text{NO}_3 \text{ used}}$$



**STEP 3** Convert moles reactant (5.68 mol)  $\rightarrow$  moles product


$$5.68 \text{ mol NH}_4\text{NO}_3 \cdot \frac{2 \text{ mol H}_2\text{O produced}}{1 \text{ mol NH}_4\text{NO}_3 \text{ used}}$$

$$= 11.4 \text{ mol H}_2\text{O produced}$$

How many moles of  $\text{N}_2\text{O}$  produced?  
Answer = **5.68 mol  $\text{N}_2\text{O}$**

MAR

MAR




**STEP 4** Convert moles product (11.4 mol)  $\rightarrow$  mass product

This is called the **THEORETICAL YIELD**

$$11.4 \text{ mol H}_2\text{O} \cdot \frac{18.02 \text{ g}}{1 \text{ mol}} = 204 \text{ g H}_2\text{O}$$

**ALWAYS FOLLOW THESE STEPS IN SOLVING STOICHIOMETRY PROBLEMS!**



**STEP 5 How much  $\text{N}_2\text{O}$  is formed?**

Total mass of reactants =  
 total mass of products  
**454 g  $\text{NH}_4\text{NO}_3 = \text{___ g } \text{N}_2\text{O} + 204 \text{ g } \text{H}_2\text{O}$**   
**mass of  $\text{N}_2\text{O} = 250. \text{ g}$  law of mass action!**

could also turn mol  $\text{NH}_4\text{NO}_3$  into mol  $\text{N}_2\text{O}$ , then grams of  $\text{N}_2\text{O}$ :

**5.68 mol  $\text{N}_2\text{O} * 44.01 \text{ g/mol} = 250. \text{ g}$**


MAR



Compound	$\text{NH}_4\text{NO}_3$	$\text{N}_2\text{O}$	$\text{H}_2\text{O}$
Initial (g)	454 g	0	0
Initial (mol)	5.68mol	0	0
Change (mol)	-5.68	+5.68	+2(5.68)
Final (mol)	0	5.68	11.4
Final (g)	0	250.	204

*Mass is conserved!*

MAR




**STEP 6 Calculate the percent yield**

We predicted a yield of 250. g of  $\text{N}_2\text{O}$ . If you isolated only 131 g of  $\text{N}_2\text{O}$ , what is the percent yield of  $\text{N}_2\text{O}$ ?

This compares the theoretical yield (250. g) and actual yield (131 g) of  $\text{N}_2\text{O}$ .

MAR




**STEP 6 Calculate the percent yield**

$$\% \text{ yield} = \frac{\text{actual yield}}{\text{theoretical yield}} \cdot 100\%$$

$$\% \text{ yield} = \frac{131 \text{ g}}{250. \text{ g}} \cdot 100\% = 52.4\%$$

MAR

**GENERAL PLAN FOR STOICHIOMETRY CALCULATIONS**




We use the "grams - moles - moles - grams" conversion often in chemistry (mass is grams)  
 Sometimes I call this the "grams - moles - moles - grams" dance lol

**Molarity in next chapter - See Stoichiometry Guide**


MAR

MAR

**PROBLEM: Using 5.00 g of  $\text{H}_2\text{O}_2$ , what mass of  $\text{O}_2$  and of  $\text{H}_2\text{O}$  can be obtained?**



**PROBLEM:** Using 5.00 g of  $\text{H}_2\text{O}_2$ , what mass of  $\text{O}_2$  and of  $\text{H}_2\text{O}$  can be obtained?



Reaction is catalyzed by  $\text{MnO}_2$

Step 1: moles of  $\text{H}_2\text{O}_2$

Step 2: use **STOICHIOMETRIC FACTOR** to calculate moles of  $\text{O}_2$

Step 3: mass of  $\text{O}_2$  (2.35 g)

Step 4: mass of  $\text{H}_2\text{O}$  (2.65 g)

*Try this problem yourself!*

MAR

## Reactions Involving a LIMITING REACTANT

In a given reaction, there is not enough of one reagent to use up the other reagent completely.

The reagent in short supply **LIMITS** the quantity of product that can be formed.



MAR

## LIMITING REACTANTS

**The Reactants**

3 hamburger patties      1/2 dozen hamburger buns      1 dozen slices of cheese

**Products**

3 cheeseburgers      3 buns left over

"Excess reactant"

1/2 dozen cheese slices left over



MAR

## LIMITING REACTANTS

**Reactants**

**Products**

$$2 \text{NO}(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2 \text{NO}_2(\text{g})$$


Limiting reactant = \_\_\_\_\_

Excess reactant = \_\_\_\_\_



MAR

## LIMITING REACTANTS




React solid Zn with 0.100 mol HCl (aq)



- Left: Balloon inflates fully, some Zn left  
\* More than enough Zn to use up the 0.100 mol HCl
- Center: Balloon inflates fully, no Zn left  
\* Right amount of each (HCl and Zn)
- Right: Balloon does not inflate fully, no Zn left.  
\* Not enough Zn to use up 0.100 mol HCl

MAR

## LIMITING REACTANTS





React solid Zn with 0.100 mol HCl (aq)



0.100 mol HCl [1 mol Zn/2 mol HCl] = 0.0500 mol Zn

	Left	Center	Right
mass Zn (g)	7.00	3.27	1.31
mol Zn	0.107	0.050	0.020
mol HCl	0.100	0.100	0.100
mol HCl/mol Zn	0.93	2.00	5.00
Lim Reactant	LR = HCl	no LR	LR = Zn


MAR

**Reaction to be Studied:**

**PROBLEM:** Mix 5.40 g of Al with 8.10 g of Cl<sub>2</sub>. How many grams of Al<sub>2</sub>Cl<sub>6</sub> can form?

MAR


*Step 1 of the Limiting Reactant problem:* Compare actual mole ratio of reactants to theoretical mole ratio.



Reactants must be in the mole ratio

$$\frac{\text{mol Cl}_2}{\text{mol Al}} = \frac{3}{2}$$

MAR


**Deciding on the Limiting Reactant**

$$\text{If } \frac{\text{mol Cl}_2}{\text{mol Al}} > \frac{3}{2}$$

then there is not enough Al to use up all the Cl<sub>2</sub>, and the limiting

reagent is **Al**

MAR

**Deciding on the Limiting Reactant**

$$\text{If } \frac{\text{mol Cl}_2}{\text{mol Al}} < \frac{3}{2}$$

then there is not enough Cl<sub>2</sub> to use up all the Al, and the limiting

reagent is **Cl<sub>2</sub>**

MAR


*Step 2 of the Limiting Reactant problem:*  
**Calculate moles of each reactant**

We have 5.40 g of Al and 8.10 g of Cl<sub>2</sub>. How much Al<sub>2</sub>Cl<sub>6</sub> can form?

$$5.40 \text{ g Al} \cdot \frac{1 \text{ mol}}{27.0 \text{ g}} = 0.200 \text{ mol Al}$$

$$8.10 \text{ g Cl}_2 \cdot \frac{1 \text{ mol}}{70.9 \text{ g}} = 0.114 \text{ mol Cl}_2$$

MAR




MAR

*Step 3 of the Limiting Reactant problem:*  
**Compare moles to find limiting reactant**


$$\frac{\text{mol Cl}_2}{\text{mol Al}} = \frac{0.114 \text{ mol}}{0.200 \text{ mol}} = 0.570$$

This  $\curvearrowright$  should be 3/2 or 1.5/1 if reactants are present in the exact stoichiometric ratio.

Limiting reagent is **Cl<sub>2</sub>**



Mix 5.40 g of Al with 8.10 g of Cl<sub>2</sub>.  
What mass of Al<sub>2</sub>Cl<sub>6</sub> can form?



MAR

MAR

**CALCULATIONS:** calculate mass of Al<sub>2</sub>Cl<sub>6</sub> expected using limiting reactant.

Step 1: Calculate moles of Al<sub>2</sub>Cl<sub>6</sub> expected using chlorine:

$$0.114 \text{ mol Cl}_2 \cdot \frac{1 \text{ mol Al}_2\text{Cl}_6}{3 \text{ mol Cl}_2} = 0.0380 \text{ mol Al}_2\text{Cl}_6$$

Step 2: Calculate mass of Al<sub>2</sub>Cl<sub>6</sub> expected based on chlorine:

$$0.0380 \text{ mol Al}_2\text{Cl}_6 \cdot \frac{266.4 \text{ g Al}_2\text{Cl}_6}{1 \text{ mol}} = 10.1 \text{ g Al}_2\text{Cl}_6$$



### Alternate Limiting Reactant Method

Calculate theoretical yield of product based on *both* reactants.

Smaller theoretical yield comes from limiting reactant, greater yield from excess reactant.

$$8.10 \text{ g Cl}_2 \cdot \frac{1 \text{ mol}}{70.9 \text{ g}} \cdot \frac{1 \text{ mol Al}_2\text{Cl}_6}{3 \text{ mol Cl}_2} \cdot \frac{266.4 \text{ g}}{1 \text{ mol}} = 10.1 \text{ g Al}_2\text{Cl}_6$$

$$5.40 \text{ g Al} \cdot \frac{1 \text{ mol}}{27.0 \text{ g}} \cdot \frac{1 \text{ mol Al}_2\text{Cl}_6}{2 \text{ mol Al}} \cdot \frac{266.4 \text{ g}}{1 \text{ mol}} = 26.6 \text{ g Al}_2\text{Cl}_6$$

10.1 g < 26.6 g, so: limiting reactant = Cl<sub>2</sub>,  
theoretical yield = 10.1 g, excess reactant = Al



MAR

MAR

How much of which reactant will remain when reaction is complete?

Cl<sub>2</sub> was the limiting reactant.

Therefore, Al was present in excess. But by how much?



First find how much Al was required based on limiting reactant (Cl<sub>2</sub>).

Then find how much Al is in excess.



### Calculating Excess Al



0.200 mol

0.114 mol = LR



$$8.10 \text{ g Cl}_2 \cdot \frac{1 \text{ mol}}{70.9 \text{ g}} \cdot \frac{2 \text{ mol Al}}{3 \text{ mol Cl}_2} \cdot \frac{26.98 \text{ g}}{1 \text{ mol}} = 2.05 \text{ g Al}$$

Excess Al = Al available - Al required  
= 5.40 g - 2.05 g  
= 3.35 g Al unused in reaction




MAR

MAR

### Using Stoichiometry to Determine a Formula


Hydrocarbons, C<sub>x</sub>H<sub>y</sub>, can be burned in oxygen to give CO<sub>2</sub> and H<sub>2</sub>O (combustion reaction).

The CO<sub>2</sub> and H<sub>2</sub>O can be collected to determine the empirical formula of the hydrocarbon.




### Using Stoichiometry to Determine a Formula

What is the empirical formula of a hydrocarbon,  $C_xH_y$ , if burning 0.115 g produces 0.379 g  $CO_2$  and 0.1035 g  $H_2O$ ?



MAR

### Using Stoichiometry to Determine a Formula




First, recognize that all C in  $CO_2$  and all H in  $H_2O$  comes from  $C_xH_y$ .

1. Calculate amount of C in  $CO_2$   
 $8.61 \times 10^{-3} \text{ mol } CO_2 \rightarrow 8.61 \times 10^{-3} \text{ mol C}$   
**1 mol C per 1 mol  $CO_2$**
2. Calculate amount of H in  $H_2O$   
 $5.744 \times 10^{-3} \text{ mol } H_2O \rightarrow 1.149 \times 10^{-2} \text{ mol H}$   
**2 mol H per 1 mol water!**

MAR

### Using Stoichiometry to Determine a Formula



Now find ratio of mol H/mol C to find values of x and y in  $C_xH_y$ .

$$\begin{aligned} &1.149 \times 10^{-2} \text{ mol H} / 8.61 \times 10^{-3} \text{ mol C} \\ &= 1.33 \text{ mol H} / 1.00 \text{ mol C} \\ &= 4 \text{ mol H} / 3 \text{ mol C} \end{aligned}$$

$$\text{Empirical formula} = C_3H_4$$


MAR

### Formulas with C, H and O

Caproic acid, the substance responsible for "dirty gym socks" smell, contains C, H and O.

Combustion analysis of 0.450 g caproic acid gives 0.418 g  $H_2O$  and 1.023 g  $CO_2$ , and the molar mass was found to be 116.2 g  $mol^{-1}$ .

What is the molecular formula of caproic acid?



**Careful:** oxygen comes from caproic acid and  $O_2$ , need special technique



MAR

### Formulas with C, H and O

Combustion analysis of 0.450 g caproic acid gives 0.418 g  $H_2O$  and 1.023 g  $CO_2$ , and the molar mass is 116.2 g  $mol^{-1}$ . What is the molecular formula?

Start with "regular" approach for mol H & mol C:

$$0.418 \text{ g } H_2O * (\text{mol}/18.02 \text{ g}) * (2 \text{ mol H/mol } H_2O) = 0.0464 \text{ mol H}$$

$$0.0464 \text{ mol H} * (1.01 \text{ g/mol H}) = 0.0469 \text{ g H}$$

$$1.023 \text{ g } CO_2 * (\text{mol}/44.01 \text{ g}) * (1 \text{ mol C/mol } CO_2) = 0.02324 \text{ mol C}$$

$$0.02324 \text{ mol C} * (12.01 \text{ g/mol C}) = 0.2791 \text{ g C}$$

Why did we convert to grams? Law of Mass Action!

MAR

### Formulas with C, H and O

0.450 g caproic acid: 0.418 g  $H_2O$  (0.0464 mol H, 0.0469 g H) and 1.023 g  $CO_2$  (0.02324 mol C, 0.2791 g C), molar mass = 116.2 g/mol. What is the molecular formula?

Realize that 0.450 g of caproic acid equals all the g C, g H and g O in the complex.

Converting mol H and mol C to grams, then subtracting from 0.450 g, gives g O in caproic acid:

$$0.450 \text{ g} - 0.0469 \text{ g} - 0.2791 \text{ g} = 0.124 \text{ g O}$$

caproic acid    g of H in acid    g of C in acid    g of O in acid

$$0.124 \text{ g O} * (\text{mol O} / 16.00 \text{ g}) = 0.00775 \text{ mol O}$$

MAR



## Formulas with C, H and O

0.450 g caproic acid: 0.418 g H<sub>2</sub>O (0.0464 mol H) and 1.023 g CO<sub>2</sub> (0.02324 mol C), molar mass = 116.2 g/mol, 0.00775 mol O. What is the molecular formula?

Now compare moles:

$C_{0.02324}H_{0.0464}O_{0.00775}$  gives  $C_3H_6O = \text{empirical formula}$

$C_3H_6O$  has a molar mass of 58.1 g/mol, which is half of the 116.2 g/mol value

Molecular Formula =  $(C_3H_6O)_2$ , or




You can now find empirical formulas based on combustion analysis (this chapter) and elemental percentages (previous chapter)!

MAR

## End of Chapter 4 Part 1

See also:

- [Chapter Four Part 1 Study Guide](#)
- [Chapter Four Part 1 Concept Guide](#)
- [Important Equations \(following this slide\)](#)
- [End of Chapter Problems \(following this slide\)](#)



MAR

Important Equations, Constants, and Handouts from this Chapter:

- be able to find the theoretical yield, actual yield, percent yield
- be able to determine the limiting reactant, excess reactant, excess reactant remaining at end of reaction
- understand how to calculate empirical formula (EF) and molecular formula (MF) using organic compounds containing oxygen

Balancing Equations:  
Reactants, Products, states of matter (s, l, g, aq), stoichiometric coefficients, Law of Conservation of Matter ("mass action")

MAR

## End of Chapter Problems: Test Yourself

See practice problem set #4 and self quizzes for balancing chemical equations examples and practice

1. What mass of Br<sub>2</sub>, in grams, is required for complete reaction with 2.56 g of Al? What mass of white, solid Al<sub>2</sub>Br<sub>6</sub> is expected? The equation:  $2 \text{Al}(s) + 3 \text{Br}_2(l) \rightarrow \text{Al}_2\text{Br}_6(s)$
2. Aluminum chloride is made by treating aluminum with chlorine:  $2 \text{Al}(s) + 3 \text{Cl}_2(g) \rightarrow 2 \text{AlCl}_3(s)$  If you begin with 2.70 g of Al and 4.05 g of Cl<sub>2</sub>, which reactant is limiting? What mass of AlCl<sub>3</sub> can be produced? What mass of the excess reactant remains when the reaction is completed?
3. Cu(NH<sub>3</sub>)<sub>4</sub>SO<sub>4</sub> is made via:  $\text{CuSO}_4(aq) + 4 \text{NH}_3(aq) \rightarrow \text{Cu}(\text{NH}_3)_4\text{SO}_4(aq)$  If you use 10.0 g of CuSO<sub>4</sub> and excess NH<sub>3</sub>, what is the theoretical yield of Cu(NH<sub>3</sub>)<sub>4</sub>SO<sub>4</sub>? If you isolate 12.6 g of Cu(NH<sub>3</sub>)<sub>4</sub>SO<sub>4</sub>, what is the percent yield of Cu(NH<sub>3</sub>)<sub>4</sub>SO<sub>4</sub>?
4. An unknown compound has the formula C<sub>x</sub>H<sub>y</sub>O<sub>z</sub>. You burn 0.0956 g of the compound and isolate 0.1356 g of CO<sub>2</sub> and 0.0833 g of H<sub>2</sub>O. What is the empirical formula of the compound? If the molar mass is 62.1 g/mol, what is the molecular formula?

MAR

## End of Chapter Problems: Answers

1. 22.7 g Br<sub>2</sub>, 25.3 g Al<sub>2</sub>Br<sub>6</sub>
2. Chlorine is limiting; 5.09 g AlCl<sub>3</sub>; 1.67 g Al remains
3. 14.3 g Cu(NH<sub>3</sub>)<sub>4</sub>SO<sub>4</sub>, 88.3%
4. EF = CH<sub>3</sub>O, MF = C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>

Be sure to view practice problem set #4 and self quizzes for balancing chemical equations examples and practice

MAR